

## **FLUCYTOSINE- flucytosine capsule**

### **Major Pharmaceuticals**

-----

### **FLUCYTOSINE Capsules USP**


**Rx only**

#### **WARNING**

**Use with extreme caution in patients with impaired renal function. Close monitoring of hematologic, renal and hepatic status of all patients is essential. These instructions should be thoroughly reviewed before administration of Flucytosine Capsules USP.**

#### **DESCRIPTION**

Flucytosine Capsules USP, an antifungal agent, is available as 250 mg and 500 mg capsules for oral administration. Each capsule also contains lactose monohydrate, colloidal silicon dioxide, talc, sodium starch glycolate, magnesium stearate and hard gelatin capsule shell which contains gelatin, purified water, black iron oxide, FD&C Blue No.1, titanium dioxide and sodium lauryl sulfate for 250 mg strength; gelatin, purified water, black iron oxide and titanium dioxide for 500 mg strength. The imprinting ink contains black iron oxide, shellac, potassium hydroxide, titanium dioxide, povidone, sodium hydroxide and FD & C Red No. 40 aluminum lake for 250 mg and 500 mg strengths. Chemically, flucytosine is 5-fluorocytosine, a fluorinated pyrimidine which is related to fluorouracil and floxuridine. It is a white to off-white crystalline powder with a molecular weight of 129.09 and the following structural formula:



#### **CLINICAL PHARMACOLOGY**

Flucytosine is rapidly and virtually completely absorbed following oral administration. Flucytosine Capsules USP is not metabolized significantly when given orally to man. Bioavailability estimated by comparing the area under the curve of serum concentrations after oral and intravenous administration showed 78% to 89% absorption of the oral dose. Peak serum concentrations of 30 to 40 µg/mL were reached within 2 hours of administration of a 2 g oral dose to normal subjects. Other studies revealed mean serum concentrations of approximately 70 to 80 µg/mL 1 to 2 hours after a dose in patients with normal renal function receiving a 6-week regimen of flucytosine (150 mg/kg/day given in divided doses every 6 hours) in combination with amphotericin B. The half-life in the majority of healthy subjects ranged between 2.4 and 4.8 hours. Flucytosine is excreted via the kidneys by means of glomerular filtration without significant tubular reabsorption. More than 90% of the total radioactivity after oral administration was recovered in the urine as intact drug. Flucytosine is deaminated (probably by gut bacteria) to 5-fluorouracil. The area under the curve (AUC) ratio of 5-fluorouracil to flucytosine is 4%. Approximately 1% of the dose is present in the urine as the α-fluoro-β-ureido-propionic acid metabolite. A small portion of the dose is excreted in the feces.

The half-life of flucytosine is prolonged in patients with renal insufficiency; the average half-life in

nephrectomized or anuric patients was 85 hours (range: 29.9 to 250 hours). A linear correlation was found between the elimination rate constant of flucytosine and creatinine clearance.

*In vitro* studies have shown that 2.9% to 4% of flucytosine is protein-bound over the range of therapeutic concentrations found in the blood. Flucytosine readily penetrates the blood-brain barrier, achieving clinically significant concentrations in cerebrospinal fluid.

### Pharmacokinetics in Pediatric Patients

Limited data are available regarding the pharmacokinetics of Flucytosine Capsules USP administered to neonatal patients being treated for systemic candidiasis. After five days of continuous therapy, median peak levels in infants were 19.6 µg/mL, 27.7 µg/mL, and 83.9 µg/mL at doses of 25 mg/kg (N=3), 50 mg/kg (N=4), and 100 mg/kg (N=3), respectively. Mean time to peak serum levels was of  $2.5 \pm 1.3$  hours, similar to that observed in adult patients. A good deal of interindividual variability was noted, which did not correlate with gestational age. Some patients had serum levels > 100 µg/mL, suggesting a need for drug level monitoring during therapy. In another study, serum concentrations were determined during flucytosine therapy in two patients (total assays performed =10). Median serum flucytosine concentrations at steady state were calculated to be  $57 \pm 10$  µg/mL (doses of 50 to 125 mg/kg/day, normalized to 25 mg/kg per dose for comparison). In three infants receiving flucytosine 25 mg/kg/day (four divided doses), a median flucytosine half-life of 7.4 hours was observed, approximately double that seen in adult patients. The concentration of flucytosine in the cerebrospinal fluid of one infant was 43 µg/mL 3 hours after a 25 mg oral dose, and ranged from 20 to 67 mg/L in another neonate receiving oral doses of 120 to 150 mg/kg/day.

## MICROBIOLOGY

### Mechanism of Action

Flucytosine is taken up by fungal organisms via the enzyme cytosine permease. Inside the fungal cell, flucytosine is rapidly converted to fluorouracil by the enzyme cytosine deaminase. Fluorouracil exerts its antifungal activity through the subsequent conversion into several active metabolites, which inhibit protein synthesis by being falsely incorporated into fungal RNA or interfere with the biosynthesis of fungal DNA through the inhibition of the enzyme thymidylate synthetase.

### Activity In Vitro

Flucytosine has been shown to be active against most strains of the following microorganisms **both *in vitro* and in clinical infections**.

*Candida albicans* and *Cryptococcus neoformans*

The following *in vitro* data are available, **but their clinical significance is unknown**.

Flucytosine exhibits *in vitro* minimum inhibitory concentrations (MIC values) of 4 µg/mL, or less against most (≥90%) strains of the following microorganisms, however, the safety and effectiveness of flucytosine in treating clinical infections due to these microorganisms have not been established in adequate and well control trials.

*Candida dubliniensis*, *Candida glabrata*, *Candida guilliermondii*, *Candida lusitanae*, *Candida parapsilosis*, *Candida tropicalis* and *Candida krusei* should be considered to be resistant to flucytosine.

*In vitro* activity of flucytosine is affected by the test conditions. It is essential to follow the approved standard method guidelines.<sup>1</sup>

### Susceptibility Testing Methods

#### ***Cryptococcus neoformans*:**

No interpretive criteria have been established for *Cryptococcus neoformans*.

#### ***Candida species*:**

**Broth Dilution Techniques:** Quantitative methods are used to determine antimicrobial minimum inhibitory concentrations (MICs). These MICs provide estimates of the susceptibility of *Candida* spp. to antimicrobial compounds. The MICs should be determined using a standardized procedure. Standardized procedures are based on a dilution method (broth)<sup>1</sup> with standardized inoculum concentrations and standardized concentrations of flucytosine powder. The MIC values should be interpreted according to the criteria in Table 1.

**Table 1. Susceptibility Interpretive Criteria for Flucytosine.**

| Antifungal Agent | Broth Dilution at 48 hours (MIC in µg/mL) |                  |               |
|------------------|---|------------------|---------------|
|                  | Susceptible (S)                           | Intermediate (I) | Resistant (R) |
| Flucytosine      | ≤4.0                                      | 8.0-16           | >32           |

A report of "Susceptible" indicates that the pathogen is likely to be inhibited if the antimicrobial compound in the blood reaches the concentration usually achievable. A report of "Intermediate" indicates that the result should be considered equivocal, and, if the microorganism is not fully susceptible to alternative, clinically feasible drugs, the test should be repeated. This category implies possible clinical applicability in body sites where the drug is physiologically concentrated or in situations where a high dosage of drug can be used. This category also provides a buffer zone which prevents small uncontrolled technical factors from causing major discrepancies in interpretation. A report of "Resistant" indicates that the pathogen is not likely to be inhibited if the antimicrobial compound in the blood reaches the concentration usually achievable; other therapy should be selected. Because of other significant host factors, *in vitro* susceptibility may not correlate with clinical outcomes.

Standardized susceptibility test procedures require the use of laboratory control microorganisms to control the technical aspects of the laboratory procedures. Standard flucytosine powder should provide the range of MIC values noted in Table 2. NOTE: Quality control microorganisms are specific strains of organisms with intrinsic biological properties relating to resistance mechanisms and their genetic expression within fungi; the specific strains used for microbiological control are not clinically significant.

**Table 2. Acceptable Quality Control Ranges for Flucytosine to be Used in Validation of Susceptibility Test Results.**

| QC Strain                              | Macrodilution<br>(MIC in µg/mL)<br>@ 48 hours | Microdilution<br>(MIC in µg/mL)<br>@ 48 hours |
|--|---|---|
| <i>Candida parapsilosis</i> ATCC 22019 | 0.12-0.5                                      | 0.12-0.5                                      |
| <i>Candida krusei</i> ATCC 6258        | 4.0-16  | 8.0-32  |

## Drug Resistance

Flucytosine resistance may arise from a mutation of an enzyme necessary for the cellular uptake or metabolism of flucytosine or from an increased synthesis of pyrimidines, which compete with the active metabolites of flucytosine (fluorinated antimetabolites). Resistance to flucytosine has been shown to develop during monotherapy after prolonged exposure to the drug.

*Candida krusei* should be considered to be resistant to flucytosine.

## Drug Combination

Antifungal synergism between flucytosine and polyene antibiotics, particularly amphotericin B has been reported *in vitro*. Flucytosine Capsules USP is usually administered in combination with amphotericin B

due to lack of cross-resistance and reported synergistic activity of both drugs.

## **INDICATIONS AND USAGE**

Flucytosine Capsules USP is indicated only in the treatment of serious infections caused by susceptible strains of *Candida* and/or *Cryptococcus*.

*Candida*: Septicemia, endocarditis and urinary system infections have been effectively treated with flucytosine. Limited trials in pulmonary infections justify the use of flucytosine.

*Cryptococcus*: Meningitis and pulmonary infections have been treated effectively. Studies in septicemias and urinary tract infections are limited, but good responses have been reported.

Flucytosine Capsules USP should be used in combination with amphotericin B for the treatment of systemic candidiasis and cryptococcosis because of the emergence of resistance to Flucytosine Capsules USP (See MICROBIOLOGY).

## **CONTRAINDICATIONS**

Flucytosine Capsules USP should not be used in patients with a known hypersensitivity to the drug.

## **WARNINGS**

Flucytosine Capsules USP must be given with extreme caution to patients with impaired renal function. Since Flucytosine Capsules USP is excreted primarily by the kidneys, renal impairment may lead to accumulation of the drug. Flucytosine Capsules USP serum concentrations should be monitored to determine the adequacy of renal excretion in such patients. Dosage adjustments should be made in patients with renal insufficiency to prevent progressive accumulation of active drug.

Flucytosine Capsules USP must be given with extreme caution to patients with bone marrow depression. Patients may be more prone to depression of bone marrow function if they: 1) have a hematologic disease, 2) are being treated with radiation or drugs which depress bone marrow, or 3) have a history of treatment with such drugs or radiation. Bone marrow toxicity can be irreversible and may lead to death in immunosuppressed patients. Frequent monitoring of hepatic function and of the hematopoietic system is indicated during therapy.

## **PRECAUTIONS**

### **General**

Before therapy with Flucytosine Capsules USP is instituted, electrolytes (because of hypokalemia) and the hematologic and renal status of the patient should be determined (see WARNINGS). Close monitoring of the patient during therapy is essential.

### **Laboratory Tests**

Since renal impairment can cause progressive accumulation of the drug, blood concentrations and kidney function should be monitored during therapy. Hematologic status (leucocyte and thrombocyte count) and liver function (alkaline phosphatase, SGOT and SGPT) should be determined at frequent intervals during treatment as indicated.

### **Drug Interactions**

Cytosine arabinoside, a cytostatic agent, has been reported to inactivate the antifungal activity of Flucytosine Capsules USP by competitive inhibition. Drugs which impair glomerular filtration may prolong the biological half-life of flucytosine.

## Drug/Laboratory Test Interactions

Measurement of serum creatinine levels should be determined by the Jaff  reaction, since Flucytosine Capsules USP does not interfere with the determination of creatinine values by this method. Most automated equipment for measurement of creatinine makes use of the Jaff  reaction.

## Carcinogenesis, Mutagenesis, Impairment of Fertility

Flucytosine has not undergone adequate animal testing to evaluate carcinogenic potential. The mutagenic potential of flucytosine was evaluated in Ames-type studies with five different mutants of *S. typhimurium* and no mutagenicity was detected in the presence or absence of activating enzymes. Flucytosine was nonmutagenic in three different repair assay systems (i.e., rec, uvr and pol).

There have been no adequate trials in animals on the effects of flucytosine on fertility or reproductive performance. The fertility and reproductive performance of the offspring (F<sub>1</sub> generation) of mice treated with 100 mg/kg/day (345 mg/M<sup>2</sup>/day or 0.059 times the human dose), 200 mg/kg/day (690 mg/M<sup>2</sup>/day or 0.118 times the human dose) or 400 mg/kg/day (1380 mg/M<sup>2</sup>/day or 0.236 times the human dose) of flucytosine on days 7 to 13 of gestation was studied; the *in utero* treatment had no adverse effect on the fertility or reproductive performance of the offspring.

## Pregnancy

### Teratogenic Effects. Pregnancy Category C

Flucytosine was shown to be teratogenic (vertebral fusions) in the rat at doses of 40 mg/kg/day (298 mg/M<sup>2</sup>/day or 0.051 times the human dose) administered on days 7 to 13 of gestation. At higher doses (700 mg/kg/day; 5208 mg/M<sup>2</sup>/day or 0.89 times the human dose administered on days 9 to 12 of gestation), cleft lip and palate and micrognathia were reported. Flucytosine was not teratogenic in rabbits up to a dose of 100 mg/kg/day (1423 mg/M<sup>2</sup>/day or 0.243 times the human dose) administered on days 6 to 18 of gestation. In mice, 400 mg/kg/day of flucytosine (1380 mg/M<sup>2</sup>/day or 0.236 times the human dose) administered on days 7 to 13 of gestation was associated with a low incidence of cleft palate that was not statistically significant. Studies in pregnant rats have shown that flucytosine injected intraperitoneally crosses the placental barrier. There are no adequate and well-controlled studies in pregnant women. Flucytosine Capsules USP should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

## Nursing Mothers

It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk and because of the potential for serious adverse reactions in nursing infants from Flucytosine Capsules USP, a decision should be made whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother.

## Pediatric Use

The efficacy and safety of Flucytosine Capsules USP have not been systematically studied in pediatric patients. A small number of neonates have been treated with 25 to 200 mg/kg/day of flucytosine, with and without the addition of amphotericin B, for systemic candidiasis. No unexpected adverse reactions were reported in these patients. It should be noted, however, that hypokalemia and acidemia were reported in one patient who received flucytosine in combination with amphotericin B, and anemia was observed in a second patient who received flucytosine alone. Transient thrombocytopenia was noted in two additional patients, one of whom also received amphotericin B.

## ADVERSE REACTIONS

The adverse reactions which have occurred during treatment with Flucytosine Capsules USP are grouped according to organ system affected.

*Cardiovascular:* Cardiac arrest, myocardial toxicity, ventricular dysfunction.

*Respiratory:* Respiratory arrest, chest pain, dyspnea.

*Dermatologic:* Rash, pruritus, urticaria, photosensitivity.

*Gastrointestinal:* Nausea, emesis, abdominal pain, diarrhea, anorexia, dry mouth, duodenal ulcer, gastrointestinal hemorrhage, acute hepatic injury including hepatic necrosis with possible fatal outcome in debilitated patients, hepatic dysfunction, jaundice, ulcerative colitis, enterocolitis, bilirubin elevation, increased hepatic enzymes.

*Genitourinary:* Azotemia, creatinine and BUN elevation, crystalluria, renal failure.

*Hematologic:* Anemia, agranulocytosis, aplastic anemia, eosinophilia, leukopenia, pancytopenia, thrombocytopenia, and fatal cases of bone marrow aplasia.

*Neurologic:* Ataxia, hearing loss, headache, paresthesia, parkinsonism, peripheral neuropathy, pyrexia, vertigo, sedation, convulsions.

*Psychiatric:* Confusion, hallucinations, psychosis.

*Miscellaneous:* Fatigue, hypoglycemia, hypokalemia, weakness, allergic reactions, Lyell's syndrome.

## **OVERDOSAGE**

There is no experience with intentional overdosage. It is reasonable to expect that overdosage may produce pronounced manifestations of the known clinical adverse reactions. Prolonged serum concentrations in excess of 100 µg/mL may be associated with an increased incidence of toxicity, especially gastrointestinal (diarrhea, nausea, vomiting), hematologic (leukopenia, thrombocytopenia) and hepatic (hepatitis).

In the management of overdosage, prompt gastric lavage or the use of an emetic is recommended. Adequate fluid intake should be maintained, by the intravenous route if necessary, since Flucytosine Capsules USP is excreted unchanged via the renal tract. The hematologic parameters should be monitored frequently; liver and kidney function should be carefully monitored. Should any abnormalities appear in any of these parameters, appropriate therapeutic measures should be instituted.

Since hemodialysis has been shown to rapidly reduce serum concentrations in anuric patients, this method may be considered in the management of overdosage.

## **DOSAGE AND ADMINISTRATION**

The usual dosage of Flucytosine Capsules USP is 50 to 150 mg/kg/day administered in divided doses at 6-hour intervals. Nausea or vomiting may be reduced or avoided if the capsules are given a few at a time over a 15-minute period. If the BUN or the serum creatinine is elevated, or if there are other signs of renal impairment, the initial dose should be at the lower level (see WARNINGS).

Flucytosine Capsules USP should be used in combination with amphotericin B for the treatment of systemic candidiasis and cryptococcosis because of the emergence of resistance to Flucytosine Capsules USP (See MICROBIOLOGY).

## **HOW SUPPLIED**

Flucytosine Capsules USP, 250 mg are size 2 hard gelatin capsule with blue opaque cap and grey opaque body imprinted with "NL 771" on the cap with black ink and "250" on the body with red ink filled with white to off white powder.

Cartons of 30 Capsules (6 capsules each blister pack x 5), NDC 0904-6834-07

Flucytosine Capsules USP, 500 mg are size 0 hard gelatin capsule with grey opaque cap and white

opaque body imprinted "NL 770" on the cap with black ink and "500" on the body with red ink, filled with white to off white powder.

Cartons of 30 Capsules (6 capsules each blister pack x 5), NDC 0904-6835-07

Store at 25°C (77°F); excursions permitted to 15°C - 30°C (59°F - 86°F).

## **REFERENCES**

1. Clinical and Laboratory Standards Institute. Reference Method for Broth Dilution Antifungal Susceptibility Testing of Yeasts; Approved Standard-Second Edition. NCCLS Document M27-A2, 2002 Volume 22, No 15, NCCLS, Wayne, PA, August 2002.

Manufactured by:

Novel Laboratories, Inc

Somerset, NJ 08873

Manufactured for:

Lupin Pharmaceuticals, Inc.

Baltimore, MD 21202

**Distributed By:**

**MAJOR® PHARMACEUTICALS**

17177 N Laurel Park Dr., Suite 233

Livonia, MI 48152

Refer to package label for Distributor's NDC Number

PI7700000201

Iss: 08/2017

## **Package/Label Display Panel**

Flucytosine Capsules, USP

250 mg

30 Capsules



## Package/Label Display Panel

Flucytosine Capsules, USP

500 mg

30 Capsules



MAJOR<sup>®</sup>  
PHARMACEUTICALS

NDC 0904-6835-07  
Unit Dose

FLUCYTOSINE  
CAPSULES, USP

500 mg

Rx only

30 CAPSULES

MAJOR<sup>®</sup>  
PHARMACEUTICALS

NDC 0904-6835-07  
Unit Dose

FLUCYTOSINE  
CAPSULES, USP

500 mg

Each capsule contains 500 mg flucytosine.

Usual Dosage: See product insert for dosage recommendations and other important prescribing information, precautions and warnings.

SEE INSERT FOUND AT  
WWW.MAJORPHARMACEUTICALS.COM

Store at 25°C (77°F); excursions permitted to 15°C - 30°C (59°F - 86°F).

Dispense in a tight, light-resistant container, with a child resistant closure, as defined in the USP/NF.

Keep this and all medication out of the reach of children. This Unit Dose package is not child resistant and is Intended for Institutional Use Only.

LBLN0017-01                      Rev. 02/19

Manufactured by:                      Manufactured by:  
Novel Laboratories, Inc.                      Lupin Pharmaceuticals, Inc.  
Somerset, NJ 08873                      Baltimore, MD 21202

Distributed by:  
MAJOR<sup>®</sup> PHARMACEUTICALS  
17177 N Laurel Park Dr., Suite 233  
Livonia, MI 48152 USA

3 09046 83507 3

|  |                         |                    |                              |
|--|-------------------------|--------------------|------------------------------|
| FLUCYTOSINE  |                         |                    |                              |
| flucytosine capsule  |                         |                    |                              |
|  |                         |                    |                              |
| Product Information  |                         |                    |                              |
| Product Type   | HUMAN PRESCRIPTION DRUG | Item Code (Source) | NDC:0904-6834(NDC:43386-771) |
| Route of Administration  | ORAL                    |                    |                              |
|  |                         |                    |                              |
|  |                         |                    |                              |
| Active Ingredient/Active Moiety                                |                         |                    |                              |
| Ingredient Name  |                         | Basis of Strength  | Strength                     |
| FLUCYTOSINE (UNII: D83282DT06) (FLUCYTOSINE - UNII:D83282DT06) |                         | FLUCYTOSINE        | 250 mg                       |
|  |                         |                    |                              |
| Inactive Ingredients   |                         |                    |                              |

| Ingredient Name  | Strength |
|--|----------|
| LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)                   |          |
| SILICON DIO XIDE (UNII: ETJ7Z6XBU4)                      |          |
| MAGNESIUM STEARATE (UNII: 70097M6I30)                    |          |
| TALC (UNII: 7SEV7J4R1U)                                  |          |
| TITANIUM DIO XIDE (UNII: 15FIX9V2JP)                     |          |
| FD&C BLUE NO. 1 (UNII: HBR47K3TBD)                       |          |
| FERROSO FERRIC O XIDE (UNII: XM0M87F357)                 |          |
| SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2) |          |
| GELATIN, UNSPECIFIED (UNII: 2G86QN327L)                  |          |
| WATER (UNII: 059QF0KO0R)                                 |          |
| SODIUM LAURYL SULFATE (UNII: 368GB514IJ)                 |          |
| SHELLAC (UNII: 46N107B7IO)                               |          |
| POTASSIUM HYDRO XIDE (UNII: WZH3C48M4T)                  |          |
| POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)                 |          |
| SODIUM HYDRO XIDE (UNII: 55X04QC32I)                     |          |
| FD&C RED NO. 40 (UNII: WZB9127XOA)                       |          |

Product Characteristics

|          |  |              |           |
|----------|--|--------------|-----------|
| Color    | GRAY (Opaque Body) , BLUE (Opaque Cap) | Score        | no score  |
| Shape    | CAPSULE                                | Size         | 18mm      |
| Flavor   |  | Imprint Code | NL771;250 |
| Contains |  |              |           |

Packaging

| # | Item Code        | Package Description                                    | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:0904-6834-07 | 30 in 1 CARTON   | 07/07/2017           |                    |
| 1 |                  | 1 in 1 BLISTER PACK; Type 0: Not a Combination Product |                      |                    |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| ANDA               | ANDA204652                               | 07/07/2017           |                    |

FLUCYTOSINE

flucytosine capsule

Product Information

|                         |                         |                    |                              |
|-------------------------|-------------------------|--------------------|------------------------------|
| Product Type            | HUMAN PRESCRIPTION DRUG | Item Code (Source) | NDC:0904-6835(NDC:43386-770) |
| Route of Administration | ORAL                    |                    |                              |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|-----------------|-------------------|----------|
|-----------------|-------------------|----------|

|  |  |  |                      |                    |
|--|--|--|----------------------|--------------------|
| FLUCYTOSINE (UNII: D83282DT06) (FLUCYTOSINE - UNII:D83282DT06) |  |  | FLUCYTOSINE          | 500 mg             |
|  |  |  |                      |                    |
| Inactive Ingredients   |  |  |                      |                    |
| Ingredient Name  |  |  | Strength             |                    |
| LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)                         |  |  |                      |                    |
| SILICON DIOXIDE (UNII: ETJ7Z6XBU4)                             |  |  |                      |                    |
| MAGNESIUM STEARATE (UNII: 70097M6I30)                          |  |  |                      |                    |
| TALC (UNII: 7SEV7J4R1U)  |  |  |                      |                    |
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP)                            |  |  |                      |                    |
| FERROSO FERRIC OXIDE (UNII: XM0M87F357)                        |  |  |                      |                    |
| SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)       |  |  |                      |                    |
| GELATIN, UNSPECIFIED (UNII: 2G86QN327L)                        |  |  |                      |                    |
| WATER (UNII: 059QF0KO0R)                                       |  |  |                      |                    |
| SHELLAC (UNII: 46N107B71O)                                     |  |  |                      |                    |
| POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)                         |  |  |                      |                    |
| POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)                       |  |  |                      |                    |
| SODIUM HYDROXIDE (UNII: 55X04QC32I)                            |  |  |                      |                    |
| FD&C RED NO. 40 (UNII: WZB9127XOA)                             |  |  |                      |                    |
|  |  |  |                      |                    |
| Product Characteristics  |  |  |                      |                    |
| Color  | WHITE (Opaque Body) , GRAY (Opaque Cap)  |  | Score                | no score           |
| Shape  | CAPSULE                                  |  | Size                 | 21mm               |
| Flavor   |  |  | Imprint Code         | NL770;500          |
| Contains   |  |  |                      |                    |
|  |  |  |                      |                    |
| Packaging  |  |  |                      |                    |
| #  | Item Code                                | Package Description                                    | Marketing Start Date | Marketing End Date |
| 1  | NDC:0904-6835-07                         | 30 in 1 CARTON   | 07/07/2017           |                    |
| 1  |  | 1 in 1 BLISTER PACK; Type 0: Not a Combination Product |                      |                    |
|  |  |  |                      |                    |
| Marketing Information  |  |  |                      |                    |
| Marketing Category   | Application Number or Monograph Citation |  | Marketing Start Date | Marketing End Date |
| ANDA   | ANDA204652                               |  | 07/07/2017           |                    |

**Labeler** - Major Pharmaceuticals (191427277)